



FOODS & FLUIDS FOR **FAST-PITCH SOFTBALL**



Success in many sports relies on each individual athlete doing their part on behalf of the team. Athletes set individual and team performance goals for the season, but rarely set nutrition goals. For example, one goal might be to arrive at practices hydrated and properly fueled in preparation to work hard. Good nutrition and hydration practices are one of several important behaviors that together can be key to successful individual performances.

Every team sport is different, and factors such as rules of play, frequency of games, length of season and position-specific requirements alter the nutritional plans. As a strength and power sport, softball players do not cover large distances but rely on short bursts of energy throughout a practice or game that can last several hours. Certain positions, such as the pitcher and catcher, engage in more frequent bursts of energy when in the field than others. Therefore, one of the greatest nutrition considerations is the consumption of adequate carbohydrate to maintain glycogen levels (energy stores) over the course of a practice or game to support the frequent high-intensity bursts of muscle activity.¹

This guide provides an overview of sports nutrition guidelines for women's fastpitch softball, which should be adapted to the individual athlete's needs. It should be noted that off-season workouts and training programs likely require different considerations, based on the nature and goals of the off-season program. For example, a softball player may have a goal to gain lean mass in the off-season, which would require a different nutrition strategy than during-season maintenance of lean mass. The recommendations within are focused on practices and games in the competitive season.

PRE-PRACTICE OR GAME FOODS & FLUIDS

Carbohydrate is the primary fuel source for muscle contraction during both high- and moderate-intensity points of the game, so it is important athletes start practices and games with enough carbohydrate stored in their bodies. Eating before a practice or game tops off the body's carbohydrate stores (called glycogen), this is especially important if the practice or game is in the morning.

The pre-event meal should be eaten 1-4 hours before exercise, contain 1-4 g/kg carbohydrate and be low in protein, fiber and fat to minimize the risk of gastrointestinal upset. The exact timing and amount of carbohydrate consumed during this time should meet the individual preferences of the athlete.^{2,4} Additionally, it is recommended that athletes drink 5-7 mL/kg of fluids with sodium approximately 4 hours prior to a training session, practice or game, and another 3-5 mL/kg about 2 hours prior if they cannot urinate or if the urine is dark.^{5,6}

Ingesting carbohydrate within the hour prior to a training session, practice or game essentially begins to meet the athlete's during-exercise fueling needs,^{2,4} and may also help the athlete decrease feelings of hunger. The amount and form of carbohydrate, such as a beverage, chew or solid food, is the individual choice of the athlete.

SUGGESTED DAILY MACRONUTRIENT INTAKE

(per kilogram of body weight)

Carbohydrate:² 5-7 g/kg/day

Protein:^{2,3} 1.2-2.0 g/kg/day



OPTIONS TO PROVIDE CARBOHYDRATE ENERGY SHORTLY BEFORE PRACTICES AND GAMES

	Serving Size	Carbohydrate	Sodium
Gatorade Energy Chews	6 chews	25 g	70 mg
Tiny Twists Pretzels	17 pretzels (28 g)	23 g	450 mg
Banana	1 medium	27 g	1 mg



PRE-PRACTICE OR GAME KEY MESSAGES

- Softball players should consume carbohydrate before a practice or game to ensure adequate carbohydrate is stored in the muscle. Carbohydrate is the primary fuel for both the high-intensity bursts of muscle contraction and prolonged muscle contractions that occur during “stop and go” activity.
- Drink 5-7 ml/kg (2.3-3.1 oz/lb) of fluid with sodium starting about 4 hours before a practice or game.

SAMPLE PRE-PRACTICE/GAME MEALS

(Examples for a 140 lb [64 kg] athlete)

Menu #1

(~4 hours prior, target ~4 g/kg carbohydrates, 256 g carbohydrate)

- Penne pasta (2 cups cooked) with 1 cup marinara sauce
- Medium piece French bread (~4 oz)
- 16 oz orange juice
- 1 cup vanilla fat-free pudding (not sugar free!) with 1 cup sliced strawberries

Totals: 1,286 calories, 256 g carbohydrate, 9 g fat, 31 g protein, 11 g fiber

Menu #2

(~3 hours prior, target ~3 g/kg carbohydrates, 192 g carbohydrate)

- Turkey sandwich
 - 3 oz fat-free deli turkey
 - Mustard/low-fat mayo (use sparingly)
 - Plain bagel
- ~40 tiny twist pretzels
- 1 large apple
- 2 fig cookies
- 20 fl oz Gatorade Endurance Formula

Totals: 890 calories, 189 g carbohydrate, 5.5 g fat, 29 g protein, 9.5 g fiber

Menu #3

(~2 hours prior, target ~2 g/kg carbohydrates, 228 g carbohydrate)

- 1 bakery-size bagel with 2 tsp low-fat cream cheese
- 1 large banana
- 20 fl oz Gatorade Endurance Formula

Totals: 550 calories, 125 g carbohydrate, 3.5 g fat, 11 g protein, 5 g fiber

DURING-PRACTICE OR GAME FOODS & FLUIDS

IMPORTANCE OF HYDRATION

Softball players spend several hours each day training, sometimes in hot and humid environments. Therefore, for both the safety and performance, paying attention to hydration is important. Athletes should be sure to drink enough fluid to prevent dehydration without overdrinking. It is generally accepted that dehydration of a ~2% or more decrease in body weight (approximately a 3 lb loss in a 150 lb athlete) may negatively affect an athlete's performance.⁵ Excessive dehydration may strain the cardiovascular system and increase body temperature, which increases the risk of heat illness. Unpublished data from GSSI conducted sweat testing revealed that mean fluid loss in female professional Softball players was 0.57 L/h (0.3-1.1 L/h) in moderate environmental conditions and players had a -1% body mass change during testing. While this volume may not sound substantial, over the course of tournament play, sweat losses over multiple hours can add up.

Answering “yes” to any of these questions may indicate inadequate hydration:

- Am I thirsty?
- Is my urine a dark yellow color (like apple juice)?
- Is my body weight noticeably lower than yesterday?

HYDRATE THE RIGHT WAY

Since practices are often longer than games, especially early in the season, it is important to develop a hydration strategy for both practices and games. To determine an athlete's sweat rate, measure body weight before and after a training session in the same environment as a competition. Also, keep track of all the fluid consumed.

A rough estimate of sweat rate can be obtained by using the following equation: **sweat rate (L/h) = (weight loss (lbs) + fluid intake (L))/exercise time (hours)**. This measurement will likely need to be made several different times for practices and competitions, especially as the weather changes. Reference the Sweat Rate Calculator on page 10.

SODIUM

Athletes sweat, and sweat contains sodium. Consuming fluid with sodium, such as in a sports drink, is important because sodium helps maintain the physiological desire to drink and helps retain the fluid consumed.^{8,9} Athletes, especially when training or competing for more than 2 hours or those who have high sweat losses, should replace both fluid and sodium during exercise.^{2,5,6} To estimate if an athlete is a “salty sweater,” look for white residue on dark-colored clothing after a training session.

TIPS FOR HYDRATION

- Know your sweat rate in the environments where you will train and compete to customize a plan to meet your unique needs. The catcher should pay particular attention to sweat rate as their equipment can hinder evaporation of sweat and body cooling, leading to potentially greater sweat fluid and electrolyte losses.
- Begin practices and games hydrated. Monitor your urine color; it should be a light yellow color (like lemonade) to indicate adequate hydration.
- Rehearse your game-day strategy during team practices and make sure you can tolerate the fluids without problems.
- Use sports drinks to provide fluid and electrolytes for hydration as well as carbohydrate for energy.

CARBOHYDRATE

Carbohydrate has been demonstrated to improve indices of performance in team sports, particularly intermittent high-intensity exercise capacity and cognitive functions like focus.^{9,10} Consuming carbohydrate during practices and games provides fuel to the muscle, brain and nervous system.¹⁰ The recommended amount of carbohydrate ingestion every hour of exercise for a team sport athlete, including softball players, is 30-60 g/h.^{1,3,12} The amount within this range may be tailored by the demands of each position and how active the athlete is in the field or on base. The form (solid, semisolid or liquid) of carbohydrate source should be determined by the preferences of the individual athlete. Athletes can take advantage of time in the dugout when their team is batting to consume fluid and carbohydrate.

SODIUM AND CARBOHYDRATE CONTENT OF GATORADE BEVERAGES

	Carbohydrate [g/12 fl oz]	Sodium [mg/12 fl oz]
Gatorade Thirst Quencher	21	160
G Zero	0	160
Gatorade Endurance Formula	22	310
Gatorlyte RTD	8	300
Gatorlyte Zero RTD	0	300
Gatorlytes powder*	0	780 (mg/ packet)

* Gatorlytes are not a beverage. They are a packet of electrolytes to be added to a 20 oz bottle of Gatorade Thirst Quencher.



EXAMPLES OF STRATEGIES TO MEET THE 30-60 G/H CARBOHYDRATE RECOMMENDATION

- 16 fl oz Gatorade Thirst Quencher
= **28 g carbohydrate**
- 32 fl oz Gatorade Thirst Quencher
= **56 g carbohydrate**
- 16 fl oz G Zero plus 12 Gatorade Prime Energy Chews
= **50 g carbohydrate**

Plan ahead to take advantage of time in the dugout to refuel.

DURING-PRACTICE OR GAME KEY MESSAGES

- Fast Pitch Softball players should determine their individual sweat rate and consume fluids with sodium to minimize body weight changes during practices and games.
- Carbohydrate intake during exercise can help maintain performance in “stop and go” activities such as Softball; athletes should aim to consume 30-60 g (120-240 calories) per hour of practice or games.
- It is possible to train the gut! If athletes are currently consuming less than the recommendations, gradually increase intake to minimize gastrointestinal issues.

POST-PRACTICE OR GAME FOODS & FLUIDS

In-season recovery nutrition should support the daily energy and hydration needs of the athlete while helping the muscles withstand the rigors of a long season. Nutrients and fluids consumed throughout the time between practices and games support recovery; highlighted here are the specific recommendations for the immediate recovery period.

Restoring the carbohydrate used from the muscle and liver during both aerobic- and anaerobic-type muscle contractions is a key focus of the post-exercise fueling needs of Fast-Pitch Softball players. When athletes have less than 8 hours between practices or games, 1.0-1.2 g/kg carbohydrate should be consumed every hour for 4 hours. When athletes have more than 8 hours between sessions, they should follow the daily carbohydrate needs for team sport athletes (5-7 g/kg/day for moderate training, or 6-10 g/kg/d during periods of heavy training) and choose carbohydrate-rich meals and snacks with protein regularly throughout the day.^{1,2}

While consuming carbohydrate for recovery will help replenish energy stores in the muscle to help the athlete be ready for the next practice or game, eating protein is important to rebuild muscle and adapt to the demands of fast-pitch softball, helping the athlete recover over the course of a long season. Athletes should consume about 20-40 g¹¹ or 0.25-0.3 g/ kg¹² of protein to start the recovery process as soon as possible after each training session, practice and game to help rebuild muscle tissue as well as adapt to the demands of training. Choose a rapidly digested, complete protein rich in the amino acid leucine, such as milk, whey, meat or eggs.^{12,13} Research is emerging on the use of plant-based proteins for recovery and muscle gain. Athletes consuming plant-based proteins should ensure they are eating a variety of foods in order to meet their essential amino acid needs to support recovery and training adaptations.¹⁴

Following practices and games, athletes should drink 20-24 oz per pound of body weight lost of fluid with sodium, to replace the amounts lost during training sessions, practices and games.^{2,5}

RECOVERY FOOD OPTIONS

	Calories	Carbohydrate [g]	Fiber [g]	Protein [g]	Fat [g]	Sodium [mg]
Option 1 Gatorade Protein Recovery Shake Water (amount based on body weight changes)	270	45	1	20	1.5	320
Option 2 Gatorade Recover Whey Protein Bar Water (amount based on body weight changes)	340-370	42-43	1-2	20	9-12	160-210
Option 3 Beef jerky (2 oz) & 10 saltine crackers Water (amount based on body weight changes)	360	28	1	21	14	1,490
Option 4 Muscle Milk 100% Whey Protein Powder mixed with water plus a banana	235	30	3	25	2	160

POST-PRACTICE OR GAME KEY MESSAGES

- Restore carbohydrate after practices and games to replace used glycogen (carbohydrate stored in the muscle and liver) and to store more glycogen as an adaptation to training.
- Athletes should consume ~20g, or 0.25-0.3 g/kg of high-quality protein as soon as possible following training sessions, practices and games to help rebuild muscle tissue.
- Rehydrate with 20-24 fl oz of fluid with sodium for every pound of body weight lost during exercise.



FEMALE SPECIFIC CONSIDERATIONS

Softball players will expend a significant amount of energy through undertaking intense training loads and competitive games. To support optimal health and performance, it is important that athletes consistently meet their energy needs, to ensure that their bodies have sufficient energy available to carry out vital physiological processes. If an athlete does not fuel their body well adequately, it can result in what is known as 'low energy availability' (LEA). Research suggests that LEA is more common among female athletes than their male counterparts.¹⁵ Energy availability with and without eating disorder risk was assessed in a study involving female athletes and performing artists. Of the athletes tested, 82.4% of the softball players presented with the greatest LEA with ED risk.¹⁶ External pressures to maintain weight may still apply to these athletes even though softball is generally not considered an aesthetic sport. Softball players need support to adopt nutrition strategies to meet their energy demands.

AN EXAMPLE: PUTTING THE SCIENCE-BASED RECOMMENDATIONS INTO PRACTICE



ATHLETE PROFILE

Name: Maya

Age: 20

Weight: 175 lbs (79.5 kg)

Type of athlete: Collegiate Softball Player

Goal: To determine a fueling strategy for games

Background: Maya is the starting center fielder for her team. She is looking for help to maintain her energy levels and focus in the 5th, 6th and 7th innings of games.

PRE-GAME

We want to make sure Maya eats adequate carbohydrate before the game to top off the stores in her muscle (called glycogen), since glycogen is an important fuel source during a softball game. Weeknight home games start at 7:00 PM and Maya needs to be to the locker room by 5pm after a full day of classes. Since Maya doesn't like to eat too close to the start of a game, she will need to eat her pre-game meal about 3 hours before game time. Aiming for ~3 g of carbohydrate per kilogram of body weight and considering her favorite foods, we designed a meal to deliver 238 g of carbohydrate. Maya can be superstitious and likes to eat the same thing before every game so she knows how her stomach will react and must eat fruit snacks before a game, so we incorporated that into her pre-game meal ritual.

In the past, Maya usually ate her favorite food, a meatball sub, chips and fruit snacks before a game. To help stay closer to her traditional food but provide more carbohydrate and less fat, we suggested homemade pizza bread, with 4 medium thickness slices of French bread, pizza sauce (1/2 cup), 2 turkey meatballs and part-skim shredded mozzarella cheese (~1/2 cup) plus a medium banana. With that, she can drink a 12-oz Gatorade Thirst Quencher and 4 oz water if desired to meet her fluid needs (397-557 mL, or 13-19 fl oz) and provide additional carbohydrate.

Maya gets fairly nervous before a game so she doesn't think about eating again but does feel like she could use a little energy at the start of the game. Her typical fruit snacks are actually not a bad choice as a carbohydrate source. We took this opportunity to incorporate her superstitious snack. She now has her fruit snacks after warm-ups while listening to her coach in the locker room.

DURING THE GAME

To determine Maya's sweat rate, we attended a practice when the team was scrimmaging to simulate the game situation as closely as possible. We weighed her before and after practice and measured her fluid intake. Based on that information, we've estimated Maya's sweat rate to be 0.8 L/h (~27 fl oz/h). Maya doesn't report any issues with cramping, nor does she typically observe salt on her compression clothing following practice, so she likely has average sodium needs. Carbohydrate intake

throughout the game is going to be important for Maya to help maintain her energy level in the later innings. It will be important for us to help Maya consume carbohydrate, particularly following the 30-60 g/hour recommendation. During games when Maya is running more in the outfield, she may want to target the higher end of the range. During games where there is less action in the outfield, she could target the lower end.

Maya plays the majority of each game, which can last up to 2 hours. Since Maya has moderate fluid needs, we suggested she try to consume one 28 oz bottle of GTQ over the course of a game, which will provide up to 28 oz of fluid and 51 g of carbohydrate total. Between the 3rd and 4th inning, she can eat a sleeve of Gatorade chews to take in an additional 25 g of carbohydrate. It is important that Maya works up to this amount of fluid and carbohydrate intake by trying it in practice. She can incorporate this strategy during games by sipping on this fluid when her team is batting. This will help her to maintain energy throughout the game.

AFTER THE GAME

Good recovery practices can help an athlete persist through a rigorous Softball season. Since Maya is a starter, we want to make sure she recovers well after each practice and game. Maya's hunger can vary after games, so we recommend she drink the Gatorade Recover Protein Shake or eat the Gatorade Recover Whey Protein Bar to get ~20 g of protein to start to rebuild muscle, carbohydrate to replace the glycogen stores in her muscles, and electrolytes to help replace sodium lost in sweat. The total amount of carbohydrate she eats at this point isn't of great importance since Maya's next practice isn't for another 24 hours and this shake will serve as a bridge to her next meal. If her team was participating in a tournament, a greater focus on carbohydrate amount would be necessary. It will be easy for her to drink the shake or eat the bar while she is cooling down or getting a short recovery work out after the game. Also, since every game is different, we recommend that she weigh herself before and after each game and drink her shake, as well as drink ~20-24 oz of water for every pound of body weight lost. When she does get to her post-game meal, it should contain high-quality protein, carbohydrate, and be lower in fiber and fat.

Any opinions or scientific interpretations expressed in this document are those of the author and do not necessarily reflect the position or policy of PepsiCo, Inc.

REFERENCES

1. Holway F & Spriet L. (2011) Sport-specific nutrition: Practical strategies for team sports. *J Sports Sci.* 29(Suppl 1):S115-125.
2. Thomas DT, Erdman KA, Burke LM (2016) American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance. *Med Sci Sport Exerc.* 48:543-568.
3. Packer et al. Variable-intensity simulated team-sport exercise increases daily protein requirements in active males (2017) *Front Nutr.* 4:1-8
4. Jeukendrup A & Killer S. (2010) The myths surrounding pre-exercise carbohydrate feeding. *Ann Nutr Metab.* 57(Suppl 2):18-25.
5. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. (2007) American College of Sports Medicine position stand: Exercise and fluid replacement. *Med Sci Sport Exerc.* 39:377-390.
6. Shirreffs S, & Sawka M. (2011) Fluid and electrolyte needs for training, competition, and recovery. *J Sport Sci.* 29 (Suppl 1): S39- 46.
7. Maughan RJ & Murray R. (2001) Sports Drinks: Basic Science and Practical Aspects, Boca Raton, FL: CRC Press. 7-8:183-224.
8. Baker LB. (2023) The Fluid Replacement process: principles of beverage reformulation for athletes. *Sport Science Exchange.* 36(244):1-8.
9. Baker LB, Rollo I, Stein KW, Jeukendrup AE. (2015) Acute effects of carbohydrate supplementation on intermittent sports performance. *Nutrients.* 7:5733-5763.
10. Burke L, Hawley J, Wong S, Jeukendrup A. (2011) Carbohydrates for training and competition. *J Sports Sci.* 29(Suppl 1):S17-27.
11. Macnaughton LS, Wardle SL, Witard OC, McGlory C, Hamilton DL, Jeromson S, Lawrence CE, Wallis GA, Tipton KD. (2016). The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein. *Physiol Rep.* 4:e12893.
12. Witard O, Jackman S, Breen L, Smith K, Selby A, Tipton K. (2014) Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr.* 99:86-95.
13. Phillips S. & Van Loon L. (2011) Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.* 29(Suppl 1):S29-38.
14. Pinckaers P, Trommelen J, Snijders T, van Loon LJC. (2021) The anabolic response to plant-based protein ingestion. *Sports Med.* 51 (suppl 1):59-74.
15. Logue DM, Madigan SM, Melin A., Delahunt E, Heinen M, Donnell SM, Corish CA. (2020) Low Energy Availability in Athletes: An Updated Narrative Review of Prevalence, Risk, Within-Day Energy Balance, Knowledge, and Impact on Sports Performance. *Nutrients.* 12:835.
16. Torres-McGehee TM, Emerson DM, Pritchett K, Moore EM, Smith AB, Uriegas NA. (2021) Energy availability with or without eating disorder risk in collegiate female athletes and performing artists. *J Athl Train.* 56(9):993-1002.



CALCULATIONS/YOUR WORKSHEET

1. BODY WEIGHT

For many calculations, you need to know your body weight in kilograms. To do this calculation:

Body weight in pounds _____ / 2.2 = kg

2. DAILY MACRONUTRIENT NEEDS

Carbohydrate:

_____ body weight (kg) * 5 g/kg = grams per day

TO

_____ body weight (kg) * 7 g/kg = grams per day

Protein:

_____ body weight (kg) * 1.2 g/kg = grams per day

TO

_____ body weight (kg) * 2.0 g/kg = grams per day

Amounts within these ranges should be determined based on the requirements of the individual sport and athlete.

3. BEFORE-EXERCISE CARBOHYDRATE NEEDS

A. Enter the time before exercise you like to eat (1-4 hours): _____ (h)

B. Enter your desired amount of carbohydrate (1-4 g/kg body weight): _____ (g)

C. Pre-exercise carbohydrate intake = _____ body weight (kg) * _____ carbohydrate amount
from line 2 (g/kg) = g carbohydrate

4. BEFORE-EXERCISE FLUID NEEDS

A. 4 hours prior to exercise:

_____ body weight (kg) * 5 mL/kg = mL

TO

_____ body weight (kg) * 7 mL/kg = mL

B. 2 hours prior to exercise (if needed):

_____ body weight (kg) * 3 mL/kg = mL

TO

_____ body weight (kg) * 5 mL/kg = mL

To convert mL to oz: _____ mL * 0.03 = _____ fluid oz



5. DURING-EXERCISE CARBOHYDRATE NEEDS

The recommendation is 30-60 g/hour, no calculation needed. Amount should be determined based on the requirements of the individual sport and athlete.

6. DURING-EXERCISE FLUID NEEDS

A. Pre-exercise weight = _____ lbs

B. Fluid consumed during exercise = _____ L

(_____ fluid oz / 33.8 = _____ L)

C. Post-exercise weight = _____ lbs

D. Weight change = Pre-exercise weight _____ lbs - Post-exercise weight _____ lbs =

E. Exercise time = _____ hours

F. Sweat rate = (Weight change _____ + Fluid intake _____ L) / _____ hours = L/h

7. POST-EXERCISE CARBOHYDRATE NEEDS (WHEN <8 HOURS RECOVERY)

Body weight _____ (kg) * 1 g/kg = g carbohydrate

TO

Body weight _____ (kg) * 1.2 g/kg = g carbohydrate

8. POST-EXERCISE FLUID NEEDS

Weight lost = Pre-exercise weight _____ lbs - Post-exercise weight _____ lbs =

Fluid needs:

_____ body weight lost * 20 oz = oz

TO

_____ body weight lost * 24 oz = oz

9. POST-EXERCISE PROTEIN NEEDS

About 20 g is appropriate for most athletes; however, to calculate your individual needs use this equation:

Body weight _____ (kg) * 0.25 g = g protein

TO

Body weight _____ (kg) * 0.3 g = g protein

