



FOODS & FLUIDS FOR **HOCKEY**



Success in many sports relies on each individual athlete doing their part on behalf of the team. Athletes set individual and team performance goals for the season, but rarely set nutrition goals. For example, one goal might be to arrive at practices hydrated and properly fueled in preparation to work hard. Good nutrition and hydration practices are one of several important behaviors that together can be key to successful individual performances.

Every team sport is different, and factors such as rules of play, frequency of games, length of season and position-specific requirements alter the nutritional plans.

Ice-hockey is a fast-paced team sport, requiring both power and endurance. Players rely on frequent short bursts of energy and must handle intense, repeated contact.¹ At the same time, players skate on average between 2.25-2.5 miles (3681-4002 meters) throughout the duration of a game depending upon position.² Therefore, one of the greatest nutrition considerations is the consumption of adequate carbohydrate to support energy needs over the course of a practice and game.¹

Additionally, players should focus on hydration since the demands of the sport, the environment and the protective clothing may increase the risk of dehydration and heat illness. Hockey players should make hydration a priority when they have two-a-day practices during training camp and when the players are wearing full helmets and pads. Even though the environment may feel cold in the arena, athletes must realize they can still dehydrate if fluid intake is not adequate.

This guide provides an overview of sports nutrition guidelines for ice-hockey, which should be adapted to individual athletes based on their position. It should be noted that off-season workouts and training programs likely require different considerations, based on the nature and goals of the off-season program. For example, an ice-hockey player may have a goal to lose fat mass and gain lean mass in the off-season, which would require a different nutrition strategy than during-season maintenance of lean mass. The recommendations below are focused on practices and games in the competitive season.



SUGGESTED DAILY MACRONUTRIENT INTAKE

(per kilogram of body weight)

Carbohydrate:³ 5-7 g/kg/day

Protein:^{3,4} 1.2-2.0 g/kg/day

PRE-PRACTICE OR GAME FOODS & FLUIDS

Eating before a practice or game tops off the body's carbohydrate stores (called glycogen), especially if the workout or competition is in the morning. Carbohydrate is the primary fuel source for muscle contraction during both high- and low-intensity points of the game, so it is important athletes start practices and games with enough carbohydrate stored in their bodies.

The pre-event meal should be eaten 1-4 hours before exercise, contain 1-4 g/kg carbohydrate and be low in protein, fiber and fat to minimize the risk of gastrointestinal upset. The exact timing and amount of carbohydrate consumed during this time should meet the individual preferences of the athlete.^{3,5} Additionally, it is recommended that athletes drink 5-7 mL/kg of fluids



with sodium approximately 4 hours prior to a workout or competition, and another 3-5 mL/kg about 2 hours prior if they cannot urinate or if the urine is dark.^{6,7}

Ingesting carbohydrate within the hour prior to training or competition essentially begins to meet the athlete's during-exercise fueling needs,^{3,5} and may also help the athlete decrease feelings of hunger. The amount and form of carbohydrate, such as a beverage, chew or solid food, is the individual choice of the athlete.

SAMPLE PRE-PRACTICE/GAME MEALS

(Examples for a 140 lb [64 kg] athlete)

Menu #1

(~4 hours prior, target ~4 g/kg carbohydrates, 256 g carbohydrate)

- Penne pasta (2 cups cooked) with 1 cup marinara sauce
- Medium piece French bread (~4 oz)
- 16 oz orange juice
- 1 cup vanilla fat-free pudding (not sugar free!) with 1 cup sliced strawberries

Totals: 1,286 calories, 256 g carbohydrate, 9 g fat, 31 g protein, 11 g fiber

Menu #2

(~3 hours prior, target ~3 g/kg carbohydrates, 192 g carbohydrate)

- Turkey sandwich
 - 3 oz fat-free deli turkey
 - Mustard/low-fat mayo (use sparingly)
 - Plain bagel
- ~40 tiny twist pretzels
- 1 large apple
- 2 fig cookies
- 20 fl oz Gatorade Endurance Formula

Totals: 890 calories, 189 g carbohydrate, 5.5 g fat, 29 g protein, 9.5 g fiber

Menu #3

(~2 hours prior, target ~2 g/kg carbohydrates, 228 g carbohydrate)

- 1 bakery-size bagel with 2 tsp low-fat cream cheese
- 1 large banana
- 20 fl oz Gatorade Endurance Formula

Totals: 550 calories, 125 g carbohydrate, 3.5 g fat, 11 g protein, 5 g fiber

OPTIONS TO PROVIDE CARBOHYDRATE ENERGY SHORTLY BEFORE TRAINING AND COMPETITION

	Serving Size	Carbohydrate	Sodium
Gatorade Energy Chews	6 chews	21 g	70 mg
Tiny Twists Pretzels	17 pretzels (28 g)	23 g	450 mg
Banana	1 medium	27 g	1 mg

PRE-PRACTICE OR GAME KEY MESSAGES

- Ice hockey players should consume carbohydrate before a practice or game to ensure adequate carbohydrate is stored in the muscle. Carbohydrate is the primary fuel for both the high-intensity bursts of muscle contraction and prolonged muscle contractions that occur during "stop and go" activity.
- Adequate fluids should be consumed about 4 hours before a practice or game.
- A nutrition plan for ice hockey players should take into account the position of the athlete and equipment.

DURING-PRACTICE OR GAME FOODS & FLUIDS

DEHYDRATION

It is generally accepted that dehydration of a ~2% or more decrease in body weight (approximately a 3 lb loss in a 150 lb athlete) may negatively affect an athlete's performance.⁷ Remember that adding pads and a helmet blocks the dissipation of heat from the body and can increase the risk of dehydration and heat illness.

Answering “yes” to any of these questions may indicate inadequate hydration:

- Am I thirsty?
- Is my urine a dark yellow color (like apple juice)?
- Is my body weight noticeably lower than yesterday?

IMPORTANCE OF HYDRATION

Hockey players spend several hours each day training, sometimes twice a day, while wearing equipment. Therefore, for both safety and performance, paying attention to hydration is important. Athletes should be sure to drink enough fluid to prevent dehydration without over-drinking. Dehydration may strain the cardiovascular system and increase body temperature, which increases the risk of heat illness. Female ice hockey players have average sweat rates with Olympic ice hockey players losing ~1L/h and varsity players losing ~0.67 L/h.⁸

HYDRATE THE RIGHT WAY

Since practices are often longer than games, especially early in the season, it is important to develop a hydration strategy for both practices and games. To determine an athlete's sweat rate, measure body weight before and after a training session in the same environment as a competition. Also, keep track of all the fluid consumed.

A rough estimate of sweat rate can be obtained by using the following equation: $\text{sweat rate (L/h)} = (\text{weight loss (lbs)} + \text{fluid intake (L)}) / \text{exercise time (hours)}$. This measurement will likely need to be made several different times for practices and competitions, especially as the weather changes. Reference the Sweat Rate Calculator on page 10.

SODIUM

Athletes sweat, and sweat contains sodium. Consuming fluid with sodium, such as in a sports drink, is important because sodium helps maintain the physiological desire to drink and helps retain the fluid consumed.^{9,10} Athletes, especially when training or competing for more than 2 hours or those who have high sweat losses, should replace both fluid and sodium during exercise.^{3,6,7} To estimate if an athlete is a “salty sweater,” look for white residue on dark-colored clothing after a training session.

TIPS FOR HYDRATION

- Know your sweat rate in the environments where you will train and compete to customize a plan to meet your unique needs.
- Begin practices and games hydrated. Monitor your urine color; it should be a light yellow color (like lemonade) to indicate adequate hydration.
- Rehearse your game-day strategy during team practices and make sure you can tolerate the fluids without problems.
- Use sports drinks to provide fluid and electrolytes for hydration as well as carbohydrate for energy.

CARBOHYDRATE

Carbohydrate has been demonstrated to improve indices of performance in team sports, particularly intermittent high-intensity exercise capacity.^{11,12,13}

Consuming carbohydrate during practices and games provides fuel to the muscle, brain and nervous system.¹³ The recommended amount of carbohydrate ingestion every hour of exercise for a team sport athlete, including ice-hockey players, is 30-60 g/h.^{1,3,13} The amount within this range may be tailored by the demands of each position and the form (solid, semisolid or liquid) should be determined by the preferences of the individual athlete.

SODIUM AND CARBOHYDRATE CONTENT OF GATORADE BEVERAGES

	Carbohydrate [g/12 fl oz]	Sodium [mg/12 fl oz]
Gatorade Thirst Quencher	21	160
G2	8	160
Gatorade Endurance Formula	22	310
G Zero	0	160
Gatorlytes powder*	0	780 (mg/ packet)
Gatorlyte RTD	8	300

* Gatorlytes are not a beverage. They are a packet of electrolytes to be added to a 20 oz bottle of Gatorade Thirst Quencher.



EXAMPLES OF STRATEGIES TO MEET THE 30-60 G/H CARBOHYDRATE RECOMMENDATION

- 16 fl oz Gatorade Thirst Quencher = 28 g carbohydrate
- 32 fl oz Gatorade Thirst Quencher = 56 g carbohydrate
- 32 fl oz G2 plus 6 Gatorade Prime Energy Chews = 42 g carbohydrate

Plan ahead to take advantage of timeouts and intermissions to refuel.

DURING-PRACTICE OR GAME KEY MESSAGES

- Ice-hockey players should determine their individual sweat rate, taking into account equipment and environment, and consume fluids with sodium to minimize body weight changes during practices and games.
- Carbohydrate intake during exercise can help maintain performance in “stop and go” activities such as hockey; athletes should aim to consume 30-60 g (120-240 calories) per hour of practice or games.
- It is possible to train the gut! If athletes are currently consuming less than the recommendations, gradually increase intake to minimize gastrointestinal issues.

POST-PRACTICE OR GAME FOODS & FLUIDS

In-season recovery nutrition should support the daily energy and hydration needs of the athlete while helping the muscles withstand the rigors of a long season. Nutrients and fluids consumed throughout the time between practices and games support recovery; highlighted here are the specific recommendations for the immediate recovery period.

Restoring the carbohydrate used from the muscle and liver during both aerobic- and anaerobic-type muscle contractions is a key focus of the post-exercise fueling needs of ice-hockey players. When athletes have less than 8 hours between practices or competitions, 1.0-1.2 g/kg carbohydrate should be consumed every hour for 4 hours. When athletes have more than 8 hours between sessions, they should follow the daily carbohydrate needs for team sport athletes (5-7 g/kg/day) for moderate training, 6-10 g/kg/d during periods of heavy training and choose carbohydrate-rich meals and snacks with some protein regularly throughout the day.^{1,3}

While consuming carbohydrate for recovery will help replenish energy stores in the muscle to help the athlete be ready for the next practice or game, eating protein is important to rebuild muscle and adapt to the demands of ice-hockey, helping the athlete recover over the course of a long season. Athletes should consume about 20-40 g¹⁴ or 0.25-0.3 g/kg¹⁵ of protein to start the recovery process as soon as possible after each training session, practice and game to help rebuild muscle tissue as well as adapt to the demands of training. Choose a rapidly digested, complete protein rich in the amino acid leucine, such as milk, whey, meat or eggs.^{15,16} Research is emerging on the use of plant-based proteins for recovery and muscle gain. Athletes consuming plant-based proteins should ensure they are eating a variety of foods in order to meet their essential amino acid needs to support recovery and training adaptations.¹⁷

Following practices and games, athletes should drink 20-24 oz per pound of body weight lost of fluid with sodium, to replace the amounts lost during training and competition.^{3,6}

RECOVERY FOOD OPTIONS

	Calories	Carbohydrate [g]	Fiber [g]	Protein [g]	Fat [g]	Sodium [mg]
Option 1 Gatorade Protein Recovery Shake Water (amount based on body weight changes)	270	45	1	20	1.5	320
Option 2 Gatorade Recover Whey Protein Bar Water (amount based on body weight changes)	340-370	42-43	1-2	20	9-12	160-210
Option 3 Beef jerky (2 oz) & 10 saltine crackers Water (amount based on body weight changes)	360	28	1	21	14	1,490
Option 4 Muscle Milk 100% Whey Protein Powder mixed with water plus a banana	235	30	3	25	2	160
Option 5 Evolve Plant-Based Protein Powder mixed with water	160	21	10	20	2.5	380

POST-PRACTICE OR GAME KEY MESSAGES

- Restore carbohydrate after practices and games to replace used glycogen (carbohydrate stored in the muscle and liver) and to store more glycogen as an adaptation to training.
- Athletes should consume ~20g, or 0.25-0.3 g/kg of high-quality protein as soon as possible following training or competition to help rebuild muscle tissue.
- Rehydrate with 20-24 fl oz of fluid with sodium for every pound of body weight lost during exercise.



AN EXAMPLE: PUTTING THE SCIENCE-BASED RECOMMENDATIONS INTO PRACTICE



ATHLETE PROFILE

Name: Claire

Age: 17

Weight: 140 lbs (63.6 kg)

Type of athlete: High School Hockey Player

Goal: To determine a fueling strategy for games

Background: Claire is a left side defender for her team and averages about 20 minutes per game. She is looking for some help to maintain energy in the third period and into overtime.

PRE-GAME

We want to make sure Claire eats adequate carbohydrate before the game to top off the stores in her muscle (called glycogen), since glycogen is an important fuel source during a hockey game. Weeknight games start at 7:00 PM and school ends at 4:00 PM. Since Claire doesn't like to eat too close to the start of a game, she will need to eat her pre-game meal about 3 hours before game time. We recommend she then follow the same timing for day games, when possible. Aiming for ~3 g of carbohydrate per kilogram of body weight and considering her favorite foods, we designed a meal to deliver 191 g of carbohydrate. Claire can be superstitious and likes to eat the same thing before every game so she knows how her stomach will react and must eat crackers before a game, so we incorporated that into her pre-game meal ritual.

In the past, Claire usually ate her favorite food, pepperoni pizza, with crackers before a game. To help stay closer to her traditional food but provide more carbohydrate and less fat, we suggested homemade pizza bread, with French bread (1/3 loaf), pizza sauce (1/4 cup) and a small amount of shredded mozzarella cheese (~1/2 cup). With that, she had a 12-oz Gatorade Thirst Quencher to meet her fluid needs (318-445 mL, or 11-15 fl oz) and provide additional carbohydrate.

Claire gets fairly nervous before a game so she doesn't think about eating again but does feel like she could use a little energy at the start of the game. Since her typical crackers are baked, they are actually not a bad choice as a carbohydrate source. We took this opportunity to incorporate her superstitious snack. She now has her crackers after warm-ups while listening to her coach in the locker room.

DURING THE GAME

To determine Claire's sweat rate, we attended a practice when the team was scrimmaging to simulate the game situation as closely as possible. We weighed her before and after practice and measured her fluid intake. Based on that information, we've estimated Claire's sweat rate to be 1.3 L/h (~44 fl oz/h), which is slightly higher than average for her position.⁸ Claire doesn't report any issues with cramping, and she doesn't typically observe salt on her compression clothing following practice, so she



likely doesn't have higher-than-average sodium needs. Carbohydrate intake throughout the game is going to be important for Claire to help maintain her energy level in the third period and into overtime, if needed. Not to mention, research shows carbohydrate intake during a simulated hockey game, as well as maintaining hydration, helps skills such as pass completion and skating speed.¹¹ Therefore, it will be important for us to help Claire consume carbohydrate, particularly following the 0-30 g/ hour recommendation.

Claire averages 20 minutes of playing time and a major junior hockey game usually lasts an hour, potentially longer if overtime is needed. Since Claire has moderately high fluid needs, we suggested she try to consume one 28 oz bottle of GTQ over the course of a game, which will provide up to 28 oz of fluid and 51 g of carbohydrate. She can get the rest of her needed fluid (~14 oz) by drinking water if desired. It is important that Claire works up to this amount of fluid and carbohydrate intake by trying it in practice. She can incorporate this strategy during games by sipping on this fluid during line changes and in between periods. This will help her to maintain energy into the third period and into overtime.

AFTER THE GAME

Good recovery practices can help an athlete persist through a grinding season like in hockey. Since Claire plays a lot of minutes, we want to make sure she recovers well after each practice and game. Claire's hunger can vary after games, so we recommend she drink the Gatorade Recover Protein Shake or eat the Gatorade Recover Whey Protein Bar to get 20 g of protein to rebuild muscle, carbohydrate to replace the glycogen stores in her muscles, and electrolytes to help replace sodium lost in sweat. The total amount of carbohydrate she eats at this point isn't of great importance since Claire's next practice isn't for another 24 hours and this shake will serve as a bridge to her next meal. It will be easy for her to drink the shake or eat the bar while she is cooling down or getting a short recovery work out after the game. Also, since every game is different, we recommend that she weigh herself before and after each game and drink her shake, as well as drink ~20-24 oz of water for every pound of body weight lost. When she does get to her post-game meal, it should contain high-quality protein, carbohydrate, and be lower in fiber and fat.

Any opinions or scientific interpretations expressed in this document are those of the author and do not necessarily reflect the position or policy of PepsiCo, Inc.

REFERENCES

1. Holway F & Spriet L. (2011) Sport-specific nutrition: Practical strategies for team sports. *J Sports Sci.* 29(Suppl 1):S115-125.
2. Douglas AS & Kennedy, CR. (2020) Tracking in-match movement demands using local positioning system in world-class men's ice hockey. *J Strength Cond Res.* 34(3): 639-646.
3. Thomas DT, Erdman KA, Burke LM (2016) American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance. *Med Sci Sport Exerc.* 48:543-568.
4. Packer et al. Variable-intensity simulated team-sport exercise increases daily protein requirements in active males (2017) *Front Nutr.* 4:1-8
5. Jeukendrup A & Killer S. (2010) The myths surrounding pre-exercise carbohydrate feeding. *Ann Nutr Metab.* 57(Suppl 2):18-25.
6. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. (2007) American College of Sports Medicine position stand: Exercise and fluid replacement. *Med Sci Sport Exerc.* 39:377-390.
7. Shirreffs S, & Sawka M. (2011) Fluid and electrolyte needs for training, competition, and recovery. *J Sport Sci.* 29 (Suppl 1): S39- 46.
8. Bigg JL, Davis Gamble AS, Vermeulen TF, Boville SM, Eskedjian GS, Palmer MS, Spriet LL. (2019) Sweat Loss and Hydration Habits of Female Olympic, Varsity and Recreational Ice Hockey Players. *Int J Sport Med.* 40(6):416-422.
9. Maughan RJ & Murray R. (2001) Sports Drinks: Basic Science and Practical Aspects, Boca Raton, FL: CRC Press. 7-8:183-224.
10. Baker LB. (2023) The Fluid Replacement process: principles of beverage reformulation for athletes. *Sport Science Exchange.* 36(244):1-8.
11. Linseman ME, Palmer MS, Sprenger HM, Spriet LL. (2014). Maintaining hydration with a carbohydrate-electrolyte solution improves performance, thermoregulation, and fatigue during an ice hockey scrimmage. *Appl physiol nutr metab.* 39(11):1214-1221.
12. Baker LB, Rollo I, Stein KW, Jeukendrup AE. (2015) Acute effects of carbohydrate supplementation on intermittent sports performance. *Nutrients.* 7:5733-5763.
13. Burke L, Hawley J, Wong S, Jeukendrup A. (2011) Carbohydrates for training and competition. *J Sports Sci.* 29(Suppl 1):S17-27.
14. Macnaughton LS, Wardle SL, Witard OC, McGlory C, Hamilton DL, Jeromson S, Lawrence CE, Wallis GA, Tipton KD. (2016). The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein. *Physiol Rep.* 4:e12893.
15. Witard O, Jackman S, Breen L, Smith K, Selby A, Tipton K. (2014) Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr.* 99:86-95.
16. Phillips S. & Van Loon L. (2011) Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.* 29(Suppl 1):S29-38.
17. Pinckaers P, Trommelen J, Snijders T, van Loon LJC. (2021) The anabolic response to plant-based protein ingestion. *Sports Med.* 51 (suppl 1):59-74.

CALCULATIONS/YOUR WORKSHEET

1. BODY WEIGHT

For many calculations, you need to know your body weight in kilograms. To do this calculation:

Body weight in pounds _____ / 2.2 = kg

2. DAILY MACRONUTRIENT NEEDS

Carbohydrate:

_____ body weight (kg) * 5 g/kg = grams per day

TO

_____ body weight (kg) * 7 g/kg = grams per day

Protein:

_____ body weight (kg) * 1.2 g/kg = grams per day

TO

_____ body weight (kg) * 2.0 g/kg = grams per day

Amounts within these ranges should be determined based on the requirements of the individual sport and athlete.

3. BEFORE-EXERCISE CARBOHYDRATE NEEDS

A. Enter the time before exercise you like to eat (1-4 hours): _____ (h)

B. Enter your desired amount of carbohydrate (1-4 g/kg body weight): _____ (g)

C. Pre-exercise carbohydrate intake = _____ body weight (kg) * _____ carbohydrate amount
from line 2 (g/kg) = g carbohydrate

4. BEFORE-EXERCISE FLUID NEEDS

A. 4 hours prior to exercise:

_____ body weight (kg) * 5 mL/kg = mL

TO

_____ body weight (kg) * 7 mL/kg = mL

B. 2 hours prior to exercise (if needed):

_____ body weight (kg) * 3 mL/kg = mL

TO

_____ body weight (kg) * 5 mL/kg = mL

To convert mL to oz: _____ mL * 0.03 = _____ fluid oz



5. DURING-EXERCISE CARBOHYDRATE NEEDS

The recommendation is 30-60 g/hour, no calculation needed. Amount should be determined based on the requirements of the individual sport and athlete.

6. DURING-EXERCISE FLUID NEEDS

A. Pre-exercise weight = _____ lbs

B. Fluid consumed during exercise = _____ L

(_____ fluid oz / 33.8 = _____ L)

C. Post-exercise weight = _____ lbs

D. Weight change = Pre-exercise weight _____ lbs - Post-exercise weight _____ lbs =

E. Exercise time = _____ hours

F. Sweat rate = (Weight change _____ + Fluid intake _____ L) / _____ hours = L/h

7. POST-EXERCISE CARBOHYDRATE NEEDS (WHEN <8 HOURS RECOVERY)

Body weight _____ (kg) * 1 g/kg = g carbohydrate

TO

Body weight _____ (kg) * 1.2 g/kg = g carbohydrate

8. POST-EXERCISE FLUID NEEDS

Weight lost = Pre-exercise weight _____ lbs - Post-exercise weight _____ lbs =

Fluid needs:

_____ body weight lost * 20 oz = oz

TO

_____ body weight lost * 24 oz = oz

9. POST-EXERCISE PROTEIN NEEDS

About 20 g is appropriate for most athletes; however, to calculate your individual needs use this equation:

Body weight _____ (kg) * 0.25 g = g protein

TO

Body weight _____ (kg) * 0.3 g = g protein

